

LocoMMotion Study Patient Selection Checklist

- Documented diagnosis of multiple myeloma according to IMWG diagnostic criteria
- Previous treatment with a PI,* an IMiD† and an anti-CD38 monoclonal antibody*
- Number of prior lines:
 - Received ≥ 3 prior lines of therapy
 - OR
 - Received ≤ 2 prior lines of therapy and double refractory to a PI and an IMiD
- ECOG performance status grade of 0 or 1
- New line of treatment will start within the next 28 days
- Evidence of progressive disease:
 - Documented evidence of progressive disease on or after the last regimen
 - OR
 - Best response to the last regimen was “stable disease” and documented evidence of progressive disease within the last 6 months
- Measurable disease as defined by any of the following:
 - Serum monoclonal paraprotein (M-protein) level ≥ 1.0 g/dL
 - OR
 - Urine monoclonal paraprotein (M-protein) level ≥ 200 mg/24 hours
 - OR
 - Light chain multiple myeloma without measurable disease in the serum or the urine (serum immunoglobulin free light chain ≥ 10 mg/dL and abnormal serum immunoglobulin kappa lambda free light chain ratio)
- ≥ 18 years
- Not pregnant and does not plan to become pregnant within the study period


ECOG = Eastern Cooperative Oncology Group; IMiD = Immunomodulatory agent; IMWG = International Myeloma Working Group; PI = Proteasome inhibitor

*PIs include bortezomib, ixazomib, carfilzomib, MLN9708, marizomib, oprozomib etc.

†IMiDs include thalidomide, lenalidomide, pomalidomide etc.

*Anti-CD38 monoclonal antibodies include daratumumab, isatuximab, TAK-079, MOR202 etc.

LocoMMotion Study Patient Selection Decision Tree



ECOG = Eastern Cooperative Oncology Group; IMiD = Immunomodulatory agent; IMWG = International Myeloma Working Group; PI = Proteasome inhibitor

*PIs include bortezomib, ixazomib, carfilzomib, MLN9708, marizomib, oprozomib etc.

†IMiDs include thalidomide, lenalidomide, pomalidomide etc.

*Anti-CD38 monoclonal antibodies include daratumumab, isatuximab, TAK-079, MOR202 etc.